

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁶ : C12Q 1/68</p>	<p>A2</p>	<p>(11) International Publication Number: WO 99/18234 (43) International Publication Date: 15 April 1999 (15.04.99)</p>
<p>(21) International Application Number: PCT/EP98/06286 (22) International Filing Date: 2 October 1998 (02.10.98) (30) Priority Data: 97/12552 8 October 1997 (08.10.97) FR (71) Applicants (for all designated States except US): SUEZ LY-ONNAISE DES EAUX [FR/FR]; 72, avenue de la Liberté, F-92753 Nanterre Cedex (FR). NORTHWESTERN UNIVERSITY [US/US]; 1801 Maple Avenue, Evanston, IL 60201 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): GUILLOT, Emmanuelle [FR/FR]; 13, avenue Charles de Gaulle, F-78230 Le Pecq (FR). URBAIN, Vincent [FR/FR]; 7, route de Sartrouville, F-78110 Le Vésinet (FR). MANEM, Jacques [FR/FR]; Thebout, Alles sur Dordogne, F-24480 Le Buisson de Cadouin (FR). RITTMANN, Bruce, E. [US/US]; Apartment H2, 728 Noyes Street, Evanston, IL 60201 (US). STAHL, David, A. [US/US]; 2119 Payne Street, Evanston, IL 60201 (US). FLAX, Jodi [US/US]; Apartment 9A, 3470 North Lake Shore Drive, Chicago, IL 60657 (US). WAGNER, Michaël [DE/DE]; Einsteinstrasse 34, D-81675 München (DE).</p>		<p>(74) Agents: ARMENGAUD, Alain et al.; Cabinet Armengaud Aîné, 3, avenue Bugeaud, F-75116 Paris (FR). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>
<p>(54) Title: MEANS FOR QUALITATIVE AND QUANTITATIVE ANALYSIS OF MICROBIAL POPULATIONS POTENTIALLY PRESENT IN A SAMPLE</p> <p>(57) Abstract</p> <p>This invention relates to means of qualitative and quantitative analysis of microbial populations potentially present in a sample. These means notably comprise the use of at least one RNA-targeted oligonucleotide probe for <i>in situ</i> hybridization in whole cells; followed by the extraction of those probes which have become hybridized by separation from their target and elution from the microbial cells; as well as the detection and measurement of said extracted probes.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE:**Means for qualitative and quantitative analysis
of microbial populations
potentially present in a sample**

5

This invention may be generally described as a means of qualitative and quantitative analysis of microbial populations potentially present in a sample. More specifically, it relates to a means of qualitative and quantitative analysis using RNA-targeted oligonucleotide probes.

10

The analysis of microbial populations potentially present is required for many types of solid and fluid samples. Some notable examples are those samples obtained from a natural or biological environment such as natural water or hot springs; samples taken from humans or animals such as blood, urine, vaginal and intestinal flora; and samples from urban, agricultural and industrial environments such as food products, industrial water, industrial effluents, municipal wastewater, industrial sludge, fermentation media, aerosols, filters or air from air conditioning systems.

15

Various laboratory techniques have been developed for the qualitative and quantitative analysis of microbial populations potentially present in a given sample.

20

One familiar technique involves a count of the microorganisms that develop after the sample (or an extract thereof) is cultured on various selective nutrient media under standard conditions. This technique is simple but entails significant risks of errors and artifacts (low specificity of morphological criteria, inability to detect viable but non-culturable microorganisms, inability to detect slow-growing microorganisms, need to maintain viability of bacteria between collection and enumeration). Moreover, this technique generally requires longer than 24 hours to yield results.

25

30

A second technique, which entails the measurement of the activity of one or more enzymes, allows a rapid quantification of populations of living microorganisms (culturable microorganisms and/or microorganisms in a viable but non-culturable form). This technique can be used, in particular, to
5 monitor a set of populations, but does not achieve very high levels of specificity or sensitivity.

A third technique using immunological probes often requires a growth step and thus requires longer than 24 hours to yield results. Moreover, it frequently lacks both sensitivity and specificity (misidentification may occur
10 due to cross-reactions).

The most recent techniques are based on the use of specific DNA probes, which are generally labeled to permit detection after hybridization with their targets. Two main categories of oligonucleotide probes have been developed: those that target DNA and those that target RNA (ribosomal RNA
15 or messenger RNA).

DNA-targeted probes, although potentially highly specific, have the drawback of low sensitivity due to the few copies of the target DNA genes in each microbial cell. Although the use of PCR (polymerase chain reaction) to amplify the target DNA sequences before detection can compensate for the
20 lack of sensitivity of the DNA probes, it has several drawbacks of its own: for example, the presence of inhibitors can lead to false-negative reactions, while carry-over or similar contamination can lead to false-positive reactions. In contrast, the use of RNA-targeted probes prevents from such drawbacks. In particular, because of the large number of copies of *rRNA* that occur
25 naturally in a microorganism (actively growing cells may contain 10^4 ribosomes, each a potential probe target), the use of *rRNA*-targeted probes does not require the amplification step, thereby overcoming the constraints and artifacts associated therewith. The advantage of targeting *rRNA* is that about 85-90 percent of the total RNA in a typical cell is *rRNA*.

30 The hybridization of RNA-targeted probes can be achieved either after cell lysis, extraction and purification of the total nucleic acids of the

sample, or *in situ* on whole cells, generally after fixation (permeabilization) of the membrane (or wall) of the microorganisms potentially present in the sample.

However, cell lysis and the ensuing extraction and purification of the nucleic acids particularly total RNA, are delicate and time-consuming manipulations that require costly apparatus, trained personnel and strict experimental conditions, notably the prevention of contamination by nucleases during the procedure. This technique further implies the use of a solid support, such as a nylon membrane, onto which the purified nucleic acids are immobilized in such a way one can discriminate between them (e.g. dot-blot, slot-blot). It most generally also implies the use of radioactive probe labels, the handling of which requires special care. The cell lysis technique for RNA hybridization is therefore ill-suited to use in routine analysis either in industry or in biological laboratories.

In situ hybridization in whole cells overcomes the need for preliminary extraction of the target nucleic acids by cellular lysis with all its associated disadvantages. The FISH (Fluorescent In Situ Hybridization) process, which employs fluorescence-labeled *rRNA* probes, is one existing *in situ* technique. This type of technique, generally involving fluorescence microscopy, provides a fast and sensitive qualitative analysis on many types of sample. Today, *rRNA*-targeted probes thus hybridized *in situ* with their target within whole cells can be quantified directly on the sample (flow cytometry, microscopy), although the method is not entirely satisfactory: quantification directly on the sample is technically costly, time-consuming, requires trained personnel and does not permit an accurate quantification of hybridized probes when the sample is complex and non-uniform (e.g. floc or aggregates formed by filamentous bacteria in sewage treatment sludge; samples containing naturally fluorescent microorganisms). As a result, the technique of *in situ* hybridization in whole cells using fluorescence-labeled oligonucleotide probes has, to date, remained an essentially qualitative technique that does not provide reliable quantitative results.

To meet the need for industrial-caliber performance on samples that can be complex and/or non-uniform, this invention provides a means for analyzing, both qualitatively and quantitatively, the microbial populations potentially present in a biological sample, said means overcoming the disadvantages of prior art techniques.

The object of this invention is, therefore, a method of qualitative and quantitative analysis of the microbial population(s) potentially present in a sample, characterized in that it comprises:

- contacting the microorganisms potentially present in said sample with at least one RNA-targeted oligonucleotide probe, hereafter called specific probe, able to target a desired microbiological population, under conditions favourable to *in situ* hybridization in whole cells,
- extracting, by separation from their target and elution outside said cells, those probes which have become hybridized,
- detecting the extracted probes and measuring the amount thereof or their respective amounts.

The present invention thus advantageously enables the extraction of said probes without destruction of said cells.

As used herein, the term "microbiological population" (or "microbiological domain") means the set of microorganisms that a given probe is able to recognize by recognition of an RNA target sequence present in each member of said set. The approach is based on oligonucleotide hybridization probes complementary to RNA sequences that are diagnostic for selected phylogenetic groups which correspond, to varying degrees, to a target region typical of a type of microorganism or a whole group of microorganisms. Any probe enabling said contacting step is appropriate for the implementation of the method according to the invention. The choice of the specific probe(s) is directly related to the analysis desired for said sample. Probes can e.g. be composed of oligonucleotide sequences that can distinguish between the primary kingdoms (eukaryotes, eubacteria, archaeobacteria) and between closely related organisms (the group of

Ammonia-oxidizing β -Proteobacteria, the genus *Nitrobacter* or *Acinetobacter* or the species *Fibrobacter intestinalis*, the species *Escherichia Coli*). Probes with finer phylogenetic resolution can be derived by using the existing collections of RNA sequences. Many examples of such RNA-targeted probes are described in the prior art such as patents or patent applications, scientific publications e.g. Los Reyes *et al.* 1997, Appli. Environ. Microbiol. Vol. 63 n°3 p.1107-1117 ; Mobarry *et al.* 1996, Appli. Environ. Microbiol. Vol.62 n°6 p.2156-2162 ; Wagner *et al.* 1994, Appli. Environ. Microbiol. Vol. 60 n°3 p.792-800 ; Kane *et al.* Appli. Environ. Microbiol. Vol. 59 n°3 p. 682-686. Other examples of such probes can also be designed by the person skilled in the art. Advantageous probes are those which target ribosomal RNA (rRNA). Examples of such advantageous probes include Nb1000 (SEQ ID N°1) and Nso 1225 (SEQ ID N°2).

The method of the invention gives particularly accurate quantitative results when the cell numbers in said sample are equal to or greater than approximately 10^3 or 10^4 cells per ml. If desired, the microorganism concentration of a liquid sample can be increased by filtration or any other technique prior to implementing the method of the invention.

In a preferred arrangement of the invention, said microorganisms potentially present in the sample are also contacted with at least one probe, hereafter called "universal probe", serving to normalize the results obtained with probes targeting specific phylogenetic groups of microorganisms ("specific probes"). The amount of a specific probe in said sample may then be expressed as a ratio of the amount of said universal probe. Such an universal probe may thus enable the expression of e.g. the specific target rRNA as a percentage of the total rRNA. Examples of such "universal probes" include probes specific for any microorganism, or probes specific for bacteria, or for eukaryotes. Such "universal probes" are well-known in the art and any of them can be used as long as it enables said contacting step, and allows the desired "specific probe" normalization. Such a "universal probe" is

used in the method according to the invention similarly as a "specific probe", and accordingly is advantageously a rRNA-targeted probe.

It may be advantageous to extract the microorganisms potentially present in said sample therefrom, in particular by centrifugation, prior to
5 proceeding with any step of the method of the invention. One reason to proceed in this manner is to remove the background noise that a sample of complex composition can generate. Another reason may be to place into solution the microorganisms potentially present in a solid, gaseous or viscous sample.

10 According to one embodiment of the invention, said contacting step is performed after the cells are made to undergo a fixation step (or permeabilization step) essential for maintaining their morphological integrity, and which makes the microorganisms potentially present in said sample permeable to short oligonucleotide probes (ca 15-25 nucleotides). This
15 fixation step allows the probes to penetrate inside the microbial cells without affecting the integrity thereof, thereby attaining their target or targets *in situ*. Where applicable, said sample is homogenized prior to said fixation step in order for said at least one probe to have access to all microbial populations potentially present in the sample.

20 Said fixation is advantageously achieved by incubating said cells in a paraformaldehyde solution that is less than 10%, preferably around 4%, for 3 to 12 hours at 4°C. This fixation procedure is more particularly adapted to Gram-negative bacterias. For certain Gram-positive bacterias, said fixation step may be achieved by incubating said cells in a 100% ethanol solution.

25 Following fixation, said cells can be recovered by e.g. centrifugation and stored until use at -20°C in a buffered solution at a pH of about 7 (PBS buffer, for example) containing approximately 50% ethanol.

In a preferred arrangement of this embodiment of the invention, said fixation is followed by a dehydration step (or drying step) prior to said
30 contacting phase. Said dehydration step can thus be carried out by placing said sample in contact with at least one ethanol solution, preferably with a

series of ethanol solutions of increasing concentration, for example by placing the sample in a 70%, 80% and then 95% ethanol solution.

Advantageously, said contacting phase is performed by placing the sample in contact with said at least one probe in the presence of a solution hereafter called "hybridization solution", which comprises a denaturing agent such as sodium dodecyl sulfate (SDS) at a concentration in a 0,001-0,1% range, preferably on the order of 0.01%; Tris-HCl, pH of about 8 at a concentration in a 0,001-0,1M range, preferably on the order of 0.02M; and a salt such as sodium chloride at a concentration in a 0,1-1,5M range, preferably on the order of 0.9M. Such a contacting is advantageously performed for an incubation time comprised between about 10 minutes and about 2 hours, and at an hybridization temperature, which is preferably the optimal temperature. For each oligonucleotide probe, the hybridization conditions (temperature; concentration of salts and denaturing agents) can be indeed optimized so as to improve the specificity of the oligonucleotide probe for the corresponding RNA sequences found in the target cells. When a plurality of oligonucleotide probes is used simultaneously, these hybridization conditions can be chosen so as to take into account the optimal conditions peculiar to every probe.

It is very advantageous for the extraction of said at least one probe to be performed following the removal of excess and unbound probe or of non-specifically associated probe material placed in contact, notably by washing with a solution hereafter called "wash solution". Such a "wash solution" advantageously comprises a denaturing agent such as sodium dodecyl sulfate (SDS) at a concentration in a 0,001-0,1% range, preferably on the order of 0.02%; Tris-HCl pH of about 8 at a concentration in a 0,001-0,1M range, preferably on the order of 0.02M, and a salt such as sodium chloride at a concentration in a 0,01-0,9M range, preferably on the order of 0.1M. The formulation of the « wash solution » (e.g. salt and denaturant nature and/or concentration) is adjusted so as to achieve the appropriate stringency; i.e. the stringency necessary to the removal of non-specifically

associated probe. Thanks to such a washing step, the extraction step will be performed only on those probes which have become effectively hybridized to the desired target(s).

According to a preferred embodiment of the invention, said extraction
5 is performed by placing the microorganisms potentially present under conditions to denature enabling the denaturation of every probe specifically associated with its target sequence, notably in the presence of a probe-target denaturing agent such as one that will separate duplex DNA/DNA or DNA/RNA, and in particular the probe - target duplex under consideration,
10 and at a temperature higher than the melting temperature of the probe under consideration, notably at a temperature of about 100°C. According to a particularly preferred embodiment of the invention, said denaturing agent is formamide. Said extraction is then performed by incubating said microorganisms in formamide at 100°C for 10 minutes using a controlled
15 temperature incubator. The supernatant may then be recovered for quantification, e.g. by centrifugation. To improve detection, said extracted probes can be concentrated notably using a Speed-Vac® prior to measuring the corresponding amount of each probe.

The detection of a target-hybridized probe and the measurement of its
20 amount thus give a qualitative and quantitative analysis of the set of target-microorganisms present in the sample. It is advantageous to perform said detection and amount measurement of the extracted probes by detection and amount measurement of a label associated with or incorporated into each of the contacted probes, such as a radioactive (^{32}P , ^{35}S , ^{125}I), chemiluminescent
25 or fluorescent label. The respective amounts of probes are then measured by quantitation of the corresponding label. It is particularly advantageous to use a fluorescent label, notably fluorescein which can be easily quantified using a fluorescence spectrophotometer.

Different probes, e.g. specific probe(s) and/or universal probe(s), can
30 be placed in separate samples, or in the same sample. In the latter case, it is

possible to distinguish each probe used from the others during the detection step, for example by giving to each one its own specific label (e.g. different fluorochromes).

The method of the invention can be applied to a variety of samples.

- 5 Samples for which an analysis using the method of the invention is of particular interest include those taken from fluids such as natural water, industrial water, industrial effluents, municipal wastewater, industrial sludge, thermal mud, food liquid or gel, fermentation medium, air, gas, aerosol; samples from a building ventilation duct, air conditioning duct; samples from
10 edible solid, soil; samples from medical apparatus; human or animal samples such as blood, urine, vaginal or intestinal flora.

- The method of the invention utilizes neither microbiological culture, nor microscopy, nor an *in vitro* amplification step (like PCR) and does not
15 require any cell lysis step. It is reproducible, simple, fast (less than 3 hours), low-cost and does not require specially trained personnel. The method of the invention offers the additional advantage of being easy to automate.

- The method of the invention notably provides a qualitative and quantitative measurement of the microbiological or sanitary status of said sample and,
20 consequently, of the product from which said sample is taken. The method of the invention can therefore advantageously be combined with an alarm function relating to the quality, safety and/or sanitary monitoring of the product from which the sample is taken, notably as part of an industrial production line.

- 25 When the threshold value or set point is exceeded, the method of the invention permits the corresponding quality, safety and/or sanitary alarm to be triggered. It also permits the automatic or feedback control of a microbiological removal or enrichment process.

- This invention also relates to the application of said method to *in vitro*
30 diagnostics of infectious diseases.

Beyond applications of the "status or condition measurement" and "alarm" types, this invention relates in particular to the application of said method for the automatic or feedback control of microbiological processes such as methane fermentation of liquid manure, treatment of organic effluents, sewage treatment processes such as activated sludge treatment; or to the automatic or feedback control of a process aimed at removing or preventing the growth of microorganisms.

Thus, the method of the invention may be advantageously applied to the detection of foam formation during the implementation of activated sludge processes and/or to the feedback control of a process aimed at removing or preventing the development of such foams.

Other features and advantages of the invention will further become apparent in the following exemplary embodiments, which are given for illustrative and non-limiting purposes.

15

EXAMPLE 1: qualitative and quantitative analysis of a sample of sewage treatment reactor effluent

a) Fixation step

Samples of effluent from sewage treatment activated sludge reactors are mixed and then washed three times using a phosphate buffer solution (PBS phosphate-buffered saline) at pH 7. The sample is then incorporated into three volumes of a 4% paraformaldehyde solution and incubated for 3 to 12 hours at 4°C. Following centrifugation the supernatant is removed and the sample is again mixed with a phosphate-buffered saline solution (PBS) at pH 7. An equal volume of ethanol is added and the sample can be stored at -20°C until use.

b) Dehydration step

The fixed sample is centrifuged after adding 1 ml of 70% ethanol over the residue and resuspending the cells. The mixture is centrifuged for 5 minutes then the supernatant is removed. This procedure is repeated with 80% ethanol and then again using 95% ethanol.

5

c) Hybridization step

A water bath is prepared at the hybridization temperature required by the probe being used (the temperature depends on the length and sequence of the probe). In the example reported here, the following probes were used:

10 Probe Nb 1000 specific to the *Nitrobacter* genus, with sequence
SEQ ID n°1: 5' TGCGACCGGTCATGG 3'

Probe Nso 1225, specific to Ammonia-oxidizing β proteobacteria, with
sequence SEQ ID n°2: 5' CGCCATTGTATTACGTGTGA 3'

Probe S Univ-1390, a universal probe for any microorganism, with
15 sequence SEQ ID n°3 : 5' GACGGGCGGTGTGTACAA 3', and

Probe S Bac338, specific for bacteria, with sequence
SEQ ID n°4: 5' GCTGCCTCCCGTAGGAGT 3'.

These probes were synthesized, purified by High Performance Liquid
Chromatography (HPLC), then fluorescein-labeled at the 5' end. They are
20 available from Operon Technologies of Alameda, California (USA) or, in
France, from the Genset company based in Paris (among others).

The cells obtained from the dehydration step are resuspended in
400 μ L of a hybridization solution comprising (for 10 mL): NaCl 5M 1.8 mL;
Tris-HCl 1M 200 μ L; SDS (sodium dodecyl sulfate) 5 μ L; distilled excipient
25 water 8 mL, for ten mL. After each probe is labeled by a fluorochrome, the
necessary quantity of each probe is added (here, 1.5 nanomoles). The cells
in the hybridization solution in contact with the probes are incubated for 10
minutes to 2 hours at the hybridization temperature. The hybridization
samples are centrifuged and supernatants are removed.

30

d) Washing step

Following hybridization, the cells are washed twice for 15 minutes each time at the hybridization temperature, in a buffered washing solution comprising, for 50 mL, NaCl 5M 1 mL; Tris-HCl 1 M 9 mL; SDS 20% 50 µL. The
5 formulation of the « washing solution » (e.g. salt and denaturant) is adjusted according to need of achieving appropriate stringency, i.e. removal of non-specifically associated probe.

e) Extraction of the fluorescence by elution

10 300 µL of formamide heated to 100°C is added to the samples obtained from the washing step, and the residue is gently resuspended. Each tube is placed in 100°C for 10 minutes, preferably using a controlled temperature incubator. Centrifuge for 10 minutes. The supernatant is recovered and stored in the dark until it can be analyzed by fluorescence spectroscopy. The
15 fluorescence is quantified using a spectrofluorometer. The amounts measured in probes Nb1000 and Nso 1225 correspond to the relative amounts of *Nitrobacter* bacteria and Ammonia-oxidizing β proteobacteria contained in the sample. These amounts are compared with those measured for universal probe S Univ-1390 and bacteria probe S Bac 338.

20 This gives a percentage ratio (the relative proportion) of the microorganisms contained in the sample, which are respectively *Nitrobacter* and Ammonia-oxidizing β proteobacteria.

It is understood that this invention is not limited to the embodiments described and illustrated herein, but covers all variants thereof.

CLAIMS

1. A method of qualitative and quantitative analysis of the microbial population(s) potentially present in a sample, characterized in that it
5 comprises:

- contacting the microorganisms potentially present in said sample with at least one RNA-targeted oligonucleotide probe, hereafter called specific probe, able to target a desired microbiological population, under conditions favourable to *in situ* hybridization in whole cells,
- 10 - extracting by separation from their target and elution outside said cells those probes which have become hybridized,
- detecting the extracted probes and measuring the amount thereof or their respective amounts.

15 2. A method according to Claim 1, further characterized in that said at least one specific probe is chosen among the group consisting of Nb 1000 (SEQ ID N°1) and Nso 1225 (SEQ ID N°2).

3. A method according to Claim 1 or 2, further characterized in that said
20 microorganisms potentially present in said sample are contacted with another probe, hereafter called universal probe, serving to normalize results obtained with probes targeting specific phylogenetic groups of microorganisms.

25 4. A method according to Claim 3, further characterized in that said universal probe is chosen among the group consisting of S Univ-1390 (SEQ ID N°3) and S Bac 338 (SEQ ID N°4).

5. A method according to any one of the preceding claims, further
30 characterized in that said specific and/or universal probe(s) is a (are) rRNA-targeted probe(s).

6. A method according to any one of the preceding Claims, further characterized in that said microorganisms potentially present in said sample are extracted from said sample, notably by centrifugation.

5

7. A method according to any one of the preceding claims, further characterized in that said contacting is performed following fixation of said cells.

10 8. A method according to claim 7, further characterized in that said fixation of the cells is achieved by incubation of said cells in a solution of less than 10% paraformaldehyde, and preferably about 4%, for 3 to 12 hours at 4°C.

15 9. A method according to claim 7 or 8, further characterized in that said fixation is followed by a dehydration step, prior to said contacting step.

10. A method according to claim 9, further characterized in that said dehydration step is performed by placing said sample in contact with at least
20 one ethanol solution, and preferably with a series of ethanol solutions of increasing concentrations.

11. A method according to any one of the preceding claims, further characterized in that said contacting is performed by placing said sample in
25 contact with said at least one probe in the presence of a solution, hereafter called hybridization solution, which notably comprises a denaturing agent such as sodium dodecyl sulfate (SDS) at a concentration in a 0,001-0,1% range, preferably on the order of 0.01%, Tris-HCl, pH of about 8 at a concentration in a 0,001-0,1 M range, preferably on the order of 0.02M; and
30 a salt such as sodium chloride at a concentration in a 0,1-1,5 M range, preferably on the order of 0.9 M .

12. A method according to any one of the preceding Claims, further characterized in that said contacting phase is performed for an incubation time of about 10 minutes to about 2 hours, and at the optimal hybridization temperature.

13. A method according to any one of the preceding claims, further characterized in that said extraction of said at least one probe is performed following removal of the excess and unbound probe or of non-specifically associated probe material placed in contact, notably by washing with a solution, hereafter called wash solution, which notably comprises a denaturing agent such as sodium dodecyl sulfate (SDS) and a salt such as sodium chloride at concentrations appropriate for achieving the stringency necessary to the removal of non-specifically associated probe.

14. A method according to any one of the preceding claims, further characterized in that said extraction is performed by placing said microorganisms potentially present under conditions enabling the denaturation of every all probe specifically associated with its target sequence, notably in the presence of an agent able to denature the probe-target duplex, and at a temperature higher than the melting temperature of the probe under consideration, notably at a temperature of approximately 100°C.

15. A method according to claim 14, further characterized in that the denaturing agent is formamide.

16. A method according to any one of the preceding claims, further characterized in that said extracted probes are concentrated, notably using

the Speed-Vac®, prior to the measurement of the amount thereof or of their respective amounts.

17. A method according to any one of the preceding claims, further
5 characterized in that said detection and amount measurement of extracted probes is performed by detection and amount measurement of a label associated or incorporated into each of the contacted probes, such as a radioactive, chemiluminescent or fluorescent label, notably such as fluorescein.

10

18. A method according to any one of the preceding claims, further characterized in that said sample is taken from fluids such as natural water, industrial water, industrial effluent, municipal wastewater, industrial sludge, thermal mud, food liquid or gel, fermentation media, air, gas, aerosol, or is a
15 sample taken from a building ventilation duct, air conditioning duct, sample of food solid, sample of soil, sample from medical apparatus, or is a human or animal sample, such as blood, urine, vaginal or intestinal flora.

19. A method according to any of the preceding claims, further
20 characterized in that it is used in combination with a process for triggering an alarm in connection with quality, safety and/or sanitary monitoring of the product from which said sample has been obtained.

20. A method according to any one of the preceding claims, further
25 characterized in that it is used in *in vitro* diagnosis of an infectious disease.

21. A method according to any one of the preceding claims, further characterized in that it is used in the automatic or feedback control of a microbiological process such as methane fermentation of liquid manure,

treatment of organic effluents, sewage treatment process such as treatment by activated sludge.

22. A method according to any of the preceding claims, further
5 characterized in that it is used in the automatic or feedback control of a process relating to the removal or prevention of the development of microorganisms.

23. A method according to any of the preceding claims, characterized in
10 that it is applied in the detection of foam formation during the implementation of activated sludge processes and/or for the feedback control of a method relating to the removal or prevention of the said foams.

LIST OF SEQUENCES

(1) General information:

(i) Applicant:

- (a) Name: Suez Lyonnaise des Eaux
- (b) Address: 72 avenue de la Liberté
- (c) City: Nanterre Cedex
- (e) Country: France
- (f) Postal code: 92753

(ii) Title of the invention: Means for qualitative and quantitative analysis of the microbial populations potentially present in a sample

(iii) Number of sequences: 4

(iv) Form readable by computer:

- (a) type of storage medium: floppy disk
- (b) computer: IBM PC compatible
- (c) operating system: PC-DOS/MS-DOS
- (d) software: PatentIn Release #1.0, Version #1.30 (OEB)

(2) Information for SEQ ID n° 1:

(i) Characteristics of the sequence:

- (a) Length: 15 base pairs
- (b) Type: nucleotide
- (c) Number of strands: single
- (d) Configuration: linear

(ii) Type of molecule: other nucleic acid

(iii) Hypothetical: yes

(iv) Antisense: no

(vii) Immediate source:

(B) Clone: Nb1000

(xi) Description of the sequence: SEQ ID n° 1:
TGCGACCGGT CATGG

(3) Information for SEQ ID n° 2:

(i) Characteristics of the sequence:

- (a) Length: 20 base pairs
- (b) Type: nucleotide

- (c) Number of strands: single
- (d) Configuration: linear
- (ii) Type of molecule: other nucleic acid
- (iii) Hypothetical: yes
- (iv) Antisense: no
- (vii) Immediate source:
(B) Clone: Nb1225
- (xi) Description of the sequence: SEQ ID n° 2:
5' CGCCATTGTA TTACGTGTGA 3'

(4) Information for SEQ ID n° 3:

- (i) Characteristics of the sequence:
 - (a) Length: 18 base pairs
 - (b) Type: nucleotide
 - (c) Number of strands: single
 - (d) Configuration: linear
- (ii) Type of molecule: other nucleic acid
- (iii) Hypothetical: yes
- (iv) Antisense: no
- (vii) Immediate source:
(B) Clone: S Univ-1390
- (xi) Description of the sequence: SEQ ID n° 3:
5' GACGGGCGGTGTGTACAA 3'

(5) Information for SEQ ID n° 4:

- (i) Characteristics of the sequence:
 - (a) Length: 18 base pairs
 - (b) Type: nucleotide
 - (c) Number of strands: single
 - (d) Configuration: linear
- (ii) Type of molecule: other nucleic acid
- (iii) Hypothetical: yes
- (iv) Antisense: no
- (vii) Immediate source:
(B) Clone: S Bac338
- (xi) Description of the sequence: SEQ ID n° 4:
5' GCTGCCTCCCGTAGGAGT 3'

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68	A3	(11) International Publication Number: WO 99/18234 (43) International Publication Date: 15 April 1999 (15.04.99)
(21) International Application Number: PCT/EP98/06286 (22) International Filing Date: 2 October 1998 (02.10.98) (30) Priority Data: 97/12552 8 October 1997 (08.10.97) FR (71) Applicants (for all designated States except US): SUEZ LY-ONNAISE DES EAUX [FR/FR]; 72, avenue de la Liberté, F-92753 Nanterre Cedex (FR). NORTHWESTERN UNIVERSITY [US/US]; 1801 Maple Avenue, Evanston, IL 60201 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): GUILLOT, Emmanuelle [FR/FR]; 13, avenue Charles de Gaulle, F-78230 Le Pecq (FR). URBAIN, Vincent [FR/FR]; 7, route de Sartrouville, F-78110 Le Vésinet (FR). MANEM, Jacques [FR/FR]; Thebout, Alles sur Dordogne, F-24480 Le Buisson de Cadouin (FR). RITTMANN, Bruce, E. [US/US]; Apartment H2, 728 Noyes Street, Evanston, IL 60201 (US). STAHL, David, A. [US/US]; 2119 Payne Street, Evanston, IL 60201 (US). FLAX, Jodi [US/US]; Apartment 9A, 3470 North Lake Shore Drive, Chicago, IL 60657 (US). WAGNER, Michaël [DE/DE]; Einsteinstrasse 34, D-81675 München (DE).	(74) Agents: ARMENGAUD, Alain et al.; Cabinet Armengaud Ainé, 3, avenue Bugeaud, F-75116 Paris (FR). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 28 September 2000 (28.09.00)	
(54) Title: MEANS FOR QUALITATIVE AND QUANTITATIVE ANALYSIS OF MICROBIAL POPULATIONS POTENTIALLY PRESENT IN A SAMPLE (57) Abstract This invention relates to means of qualitative and quantitative analysis of microbial populations potentially present in a sample. These means notably comprise the use of at least one RNA-targeted oligonucleotide probe for <i>in situ</i> hybridization in whole cells; followed by the extraction of those probes which have become hybridized by separation from their target and elution from the microbial cells; as well as the detection and measurement of said extracted probes.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

International Application No
PCT/EP 98/06286

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C120

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WAGNER M ET AL.: "In situ identification of ammonia-oxidizing bacteria" SYSTEMATIC AND APPLIED MICROBIOLOGY, vol. 18, 1995, pages 251-264, XP002068767 see the whole document</p> <p style="text-align: center;">---</p>	<p>1,3, 5-11, 13-23</p>
X	<p>DE LOS REYES ET AL.: "Group-specific small-subunit rRNA hybridization probes to characterize filamentous foaming in activated sludge systems" APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 63, no. 3, 1997, pages 1107-1117, XP002068768 cited in the application see the whole document</p> <p style="text-align: center;">---</p>	<p>1-11,13, 16-18, 21-23</p>

-/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention.

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

24 March 1999

Date of mailing of the international search report

09/04/1999

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Knehr, M

INTERNATIONAL SEARCH REPORT

Int. Application No.

PCT/EP 98/06286

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MANZ W ET AL: "IN SITU CHARACTERIZATION OF THE MICROBIAL CONSORTIA ACTIVE IN TWO WASTEWATER TREATMENT PLANTS" WATER RESEARCH, vol. 28, no. 8, 1 August 1994, pages 1715-1723, XP000446344 see the whole document ---	1,3, 5-11, 16-19, 21-23
X	WAGNER M ET AL.: "Probing activated sludge with oligonucleotides specific for proteobacteria: Inadequacy of culture-dependent methods for describing microbial community structure" APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 59, no. 5, 1993, pages 1520-1525, XP002068769 see the whole document ---	1,3, 5-10,17, 18,21-23
X	MOBARRY B K ET AL.: "Phylogenetic probes for analyzing abundance and spatial organization of nitrifying bacteria" APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 62, no. 6, 1996, pages 2156-2162, XP002068770 cited in the application see abstract see page 2156, column 1, paragraph 1 - column 2, paragraph 3 see page 2159, column 2, paragraph 2 - page 2161, column 2, paragraph 3; figures 1,2; tables 1,3 ---	1-3,5, 17-19, 21-23
X	WAGNER M ET AL.: "Development of an rRNA-targeted oligonucleotide probe specific for the genus Acinobacter and its application for in situ monitoring in activated sludge" APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 60, no. 3, 1994, pages 792-800, XP002097846 see the whole document ---	1,5,7,8, 14,15, 17,18
X	MOBARRY B K ET AL.: "Phylogenetic probes for analyzing abundance and spatial organization of nitrifying bacteria" APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 62, no. 6, 1997, page 815 XP002068771 cited in the application see page 815, line 1 - line 2 ---	1,2
X	US 5 426 025 A (REEVES ROBERT H ET AL) 20 June 1995 * spécialement colonne 5, ligne 8 à colonne 6, ligne 14 * see the whole document ---	1,3,5, 17,18,20

INTERNATIONAL SEARCH REPORT

In. ational Application No
PCT/EP 98/06286

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	LEMMER (M) H ET AL: "Denitrification in a methanol-fed fixed-bed reactor. Part 2: composition and ecology of the bacterial community in the biofilms" WATER RESEARCH, vol. 31, no. 8, August 1997, page 1903-1908 XP004081404 see abstract see page 1903, column 1, paragraph 1 - page 1904, column 1, paragraph 1 -----	1,18,19, 21-23
X	WO 88 03957 A (GEN PROBE INC) 2 June 1988 see abstract; claim 219 -----	1,3-5
X	WO 96 19585 A (HEIDELBERG REPATRIATION HOSPIT ;GUERTLER VOLKER (AU)) 27 June 1996 * page 29, tableau 4 * -----	1,3-5

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. National Application No

PCT/EP 98/06286

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5426025	A	20-06-1995	US 5607835 A	04-03-1997
WO 8803957	A	02-06-1988	AT 163680 T	15-03-1998
			AU 616646 B	07-11-1991
			AU 1041988 A	16-06-1988
			CA 1339871 A	19-05-1998
			DE 3752172 D	09-04-1998
			DE 3752172 T	02-07-1998
			DK 413788 A	23-09-1988
			EP 0272009 A	22-06-1988
			ES 2112824 T	16-04-1998
			FI 883482 A	22-07-1988
			JP 10042880 A	17-02-1998
			JP 1503356 T	16-11-1989
			KR 9511719 B	09-10-1995
			PT 86204 B	07-11-1990
			US 5541308 A	30-07-1996
			US 5595874 A	21-01-1997
			US 5547842 A	20-08-1996
			US 5593841 A	14-01-1997
			US 5683876 A	04-11-1997
			US 5677127 A	14-10-1997
			US 5677128 A	14-10-1997
			US 5677129 A	14-10-1997
			US 5827651 A	27-10-1998
			US 5693468 A	02-12-1997
			US 5691149 A	25-11-1997
			US 5693469 A	02-12-1997
			US 5679520 A	21-10-1997
			US 5714321 A	03-02-1998
			US 5674684 A	07-10-1997
			US 5840488 A	24-11-1998
WO 9619585	A	27-06-1996	AU 1307595 A	10-07-1996